

Magnetic Resonance chromatography (NMR) to quantify the formation of carbonic acid and carbamate. The CO₂ capture experiments were carried out over a 24-hour period. There was a drop in the pH of glycine and sarcosine solutions after 24 hours at both concentrations and methods. The decline in pH is due to increased levels of carbamate and carbonate. In proton NMR analysis, the intensity of the proton at 3.8 decreases, while the intensity at

AIChE Meetings

2022

- May 2-5 [2022 Synthetic Biology: Evolution, Engineer and Design \(SEED\)](#)
Arlington, VA
- June 1-3 [Advanced Manufacturing and Processing Conference](#)
Bethesda, MD
- June 7-9 [Process Development Symposium](#)
Philadelphia, PA
- June 26-28 [NDEW-ChE: National Diversity Equity Workshop for Chemical Engineering Academic Leaders](#)
Baltimore, MD

National AIChE Contact Info

AIChE Customer Service Center:
(US/Canada): 1-800-AIChemE
(1-800-242-4363)
(International): 203/702-7660
E-mail: CustomerService@aiche.org
Member Services: 800.242.4363
Career Services: 646.495.1330
Awards & Honors: 646.495.1317
Address: 120 Wall Street, 23rd

MEETING SCHEDULE

The Rocky Mountain Local Section (RMLS) of AIChE generally meets the second or third Tuesday of every month, September through May.